Author index

Alejo-Blanco, R., see Kawamoto, F., 101
Alexander, K., Parail, A.C. and Parsons, M.
An allele of *Trypanosoma brucei* cytoplasmic phosphoglycerate kinase is a mosaic of other alleles and genes (*Short Communication*), 293
Aline Jr., R.F., see Smiley, B.L., 143
Arnold, H-H., see Ragge, K., 93
Åslund, L., see Henriksson, J., 213
Avila, H., Goncalves, A.M., Nehme, N.S., Morel, C.M. and Simpson, L.
Schizodeme analysis of *Trypanosoma cruzi* stocks from South and Central America by analysis of polymerase chain reaction-amplified minicircle variable region sequences, 175

Baack, B.R., see Vander Jagt, D.L., 277
Barrett, J., see Brophy, P.M., 205
Battistutta, D., see Saul, A., 55
Beck, D.J., see Lew, A.M., 153
Beverley, S.M. and Coburn, C.M.
Recurrent de novo appearance of small linear DNAs in *Leishmania major* and relationship to extrachromosomal DNAs in other species, 133
Bobek, L.A., see Henkle, K.J., 69
Borst, P., see Crozatier, M., 1
Bosch, M., see Gonzalez Clemente, M. del C., 289
Bossio, J.C., see Gonzalez Clemente, M. del C., 289
Brodin, T.N., see Sacks, D.L., 225
Brophy, P.M. and Barrett, J.
Strategies for detoxification of aldehydic products of lipid peroxidation in helminths, 205
Bzik, D.J., see Tanaka, M., 83

Cain, G.D., see Henkle, K.J., 69
Campos, N.M., see Vander Jagt, D.L., 277
Carr, D., see Thomas, A.W., 285
Cazzulo, J.J., see Henriksson, J., 215
Coburn, C.M., see Beverley, S.M., 133
Conrad, J.T., see Nash, T.E., 125
Cook, G.A., see Henkle, K.J., 69
Coombs, G.H., see Robertson, C.D., 269
Cowman, A.F. and Lew, A.M.
Chromosomal rearrangements and point mutations in the DHFR-TS gene of *Plasmodium chabaudi* under antifolate selection, 21
Crozatier, M., Van der Ploeg, L.H.T., Johnson, P.J., Gommers-Ampt, J. and Borst, P.
Structure of a telomeric expression site for variant specific surface antigens in *Trypanosoma brucei*, 1

Dietrich, P., Pilar Dussan, M. del, Floeter-Winter, L.M., Trebilcock Affonso, M.H., Plessman Camargo, E. and Soares, M.B.
Restriction fragment length polymorphisms in the ribosomal gene spacers of *Trypanosoma cruzi* and *Trypanosoma conorhini*, 13
Docampo, R., see Vercesi, A.E., 119
Donelson, J.E., see Henkle, K.J., 69

Edwards, M.R., see Grote, R., 109
Engman, D.M., see Henkle, K.J., 69

Ferrero, M.A.G., see Gonzalez Clemente, M. del C., 289
Fetterer, R.H., see Rhoads, M.L., 155
Fleck, S.L., see Kawamoto, F., 101
Floeter-Winter, L.M., see Dietrich, P., 13
Foster, L.A., see Henkle, K.J., 69
Franke de Cazzulo, B.M., see Henriksson, J., 213
Frasch, A.C.C., see Henriksson, J., 213

Gadelha, F.R., see Vercesi, A.E., 119
Gamble, H.R., see Zarlenga, D.S., 165
Garside, L., see Gibson, W., 45
Gibson, W. and Garside, L.
Kinetoplast DNA minicircles are inherited from both parents in genetic hybrids of *Trypanosoma brucei*, 45
Gommers-Ampt, J., see Crozatier, M., 1
Goncalves, A.M., see Avila, H., 175
Gonzalez Clemente, M. del C., Ferrero, M.A.G., Bossio, J.C., Bosch, M. and Parodi, A.J.
The structure of a galactofuranose-containing oligosaccharide isolated from glycoproteins of the trypanosomatid *Herpetomonas samuelpessoai* (*Short Communication*), 289
Gregory, W.F., see Selkirk, M.E., 31
Grote, R., Edwards, M.R., Norton, R.S. and O'Sullivan, W.J. A ^{31}P nuclear magnetic resonance study of *Crithidia luciliae*, 109
Gu, H.M., see Tanaka, M., 83

Henkle, K.J., Cook, G.A., Foster, L.A., Engman, D.M., Bobek, L.A., Cain, G.D. and Donelson, J.E.
The gene family encoding eggshell proteins of *Schistosoma japonicum*, 69
Henriksson, J., Åslund, L., Macina, R.A., Franke de Cazzulo, B.M., Cazzulo, J.J., Frasch, A.C.C. and Pettersson, U.
Chromosomal localisation of seven cloned antigen genes provides evidence of diploidy and further demonstration of karyotype variability in *Trypanosoma cruzi*, 213
Herwaldt, B.L., Schlesinger, P.H. and Krogstad, D.J.
Accumulation of chloroquine by membrane preparations from *Plasmodium falciparum*, 257
Holder, A., see Keen, J., 241
Howard, R.F.
The lower-molecular-weight protein complex (R_1) of the *Plasmodium falciparum* rhoptries lacks the glycolytic enzyme aldolase, 235
Hundt, E., see Ragge, K., 93
Hunsaker, L.A., see Vander Jagt, D.L., 277

Inselburg, J., see Tanaka, M., 83

Jenkins, R.E., see Selkirk, M.E., 31
Jensen, J., see Petersen, C., 187
Johnson, P.J., see Crozatier, M., 1

Kawamoto, F., Alejo-Blanco, R., Fleck, S.L., Kawamoto, Y. and Sinden, R.E.
Possible roles of Ca^{2+} and cGMP as mediators of the exflagellation of *Plasmodium berghei* and *Plasmodium falciparum*, 101
Kawamoto, Y., see Kawamoto, F., 101
Keen, J., Holder, A., Playfair, J., Lockyer, M. and Lewis, A.
Identification of the gene for a *Plasmodium yoelii* rhoptry protein. Multiple copies in the parasite genome, 241
Knapp, B., see Ragge, K., 93
Krogstad, D.J., see Herwaldt, B.L., 257

Leech, J., see Petersen, C., 187
Lew, A.M., see Cowman, A.F., 21
Lew, A.M. and Beck, D.J.
The epitope of a protective monoclonal antibody occurs in a region of microheterogeneity in *Plasmodium chabaudi* (*Short Communication*), 153
Lewis, A., see Keen, J., 239
Li, W-B., see Tanaka, M., 83
Lima, S.A., see Vercesi, A.E., 119
Lingelbach, K., see Ragge, K., 93
Lloyd, D., see Paget, T.A., 63
Lockyer, M., see Keen, J., 241

Macedo, D.V., see Vercesi, A.E., 119
Macina, R.A., see Henriksson, J., 213
Maizels, R.M., see Selkirk, M.E., 31
Merritt, Jr., J.W., see Nash, T.E., 125
Morel, C.M., see Avila, H., 175
Myler, P.J., see Smiley, B.L., 143

Nash, T.E., Conrad, J.T. and Merritt Jr., J.W.
Variant specific epitopes of *Giardia lamblia*, 125
Nehme, N.S., see Avila, H., 175
Nelson, R., see Petersen, C., 189
Nielsen, B., see Parsons, M., 197
Norton, R.S., see Grote, R., 109

O'Sullivan, W.J., see Grote, R., 109

Paget, T.A., Raynor, M.H., Shipp, D.W.E. and Lloyd, D.
Giardia lamblia produces alanine anaerobically but not in the presence of oxygen, 63
Parail, A.C., see Alexander, K., 293
Parodi, A.J., see Gonzalez Clemente, M. del C., 289
Parsons, M. and Nielsen, B.
Active transport of 2-deoxy-D-glucose in *Trypanosoma brucei* procyclic forms, 197
Parsons, M., see Alexander, K., 293
Pedraza, G., see Shapira, M., 247
Petersen, C., Nelson, R., Leech, J., Jensen, J., Wollish, W. and Scherf, A.
The gene product of the *Plasmodium falciparum* 11.1 locus is a protein larger than one megadalton, 189
Pettersson, U., see Henriksson, J., 213
Pilar Dussan, M. del, see Dietrich, P., 13
Playfair, J., see Keen, J., 241
Plessman Camargo, E., see Dietrich, P., 13

Ragge, K., Arnold, H-H., Tummler, M., Knapp, B., Hundt, E. and Lingelbach, K.
In vitro biosynthesis and membrane translocation of the serine rich protein of *Plasmodium falciparum*, 93
Raynor, M.H., see Paget, T.A., 63
Rhoads, M.L. and Fetterer, R.H.
Biochemical and immunochemical characterization of ^{125}I-labeled cuticle components of *Haemonchus contortus*, 155
Robertson, C.D. and Coombs, G.H.
Characterisation of three groups of cysteine proteinase in the amastigotes of *Leishmania mexicana mexicana*, 269

Sacks, D.L., Brodin, T.N. and Turco, S.J.
Developmental modification of the lipophosphoglycan from *Leishmania major* promastigotes during metacyclogenesis, 225
Saul, A. and Battistutta, D. Analysis of the sequences flanking the translational start sites of *Plasmodium falciparum*, 55
Scherf, A., see Petersen, C., 189
Schlesinger, P.H., see Herwaldt, B.L., 257
Selkirk, M.E., Gregory, W.F., Yazdanbakhsh, M., Jenkins, R.E. and Maizels, R.M.
Cuticular localisation and turnover of the major surface glycoprotein (gp29) of adult *Brugia malayi*, 31
Shapira, M. and Pedraza, G.
Sequence analysis and transcriptional activation of heat shock protein 83 of *Leishmania mexicana amazonensis*, 247
Shipp, D.W.E., see Paget, T.A., 63
Simpson, L., see Avila, H., 175
Sinden, R.E., see Kawamoto, F., 101
Smiley, B.L., Aline, Jr., R.F., Myler, P.J. and Stuart, K.
A retroposon in the 5′ flank of a *Trypanosoma brucei* VSG gene lacks insertional terminal repeats, 143
Soares, M.B., see Dietrich, P., 13
Stuart, K., see Smiley, B.L., 143

Tanaka, M., Gu, H.-M., Bzik, D.J., Li, W-B. and Inselburg, J.
Mutant dihydrofolate reductase-thymidylate synthase genes in pyrimethamine-resistant *Plasmodium falciparum* with polymorphic chromosome duplications, 83
Thomas, A.W., Waters, A.P. and Carr, D.
Analysis of variation in PF83, an erythocytic merozoite vaccine candidate antigen of *Plasmodium falciparum* (*Short Communication*), 285
Trebilcock Affonso, M.H., see Dietrich, P., 13
Tummler, M., see Ragge, K., 93
Turco, S.J., see Sacks, D.L., 225

Van der Ploeg, L.H.T., see Crozatier, M., 1
Vander Jagt, D.L., Hunsaker, L.A., Campos, N.M. and Baack, B.R.
D-Lactate production in erythrocytes infected with *Plasmodium falciparum*, 277
Vercesi, A.E., Macedo, D.V., Lima, S.A., Gadelha, F.R. and Docampo, R.
Ca^{2+} transport in digitonin-permeabilized trypanosomatids, 119

Waters, A.P., see Thomas, A.W., 285
Wollish, W., see Petersen, C., 189

Yazdanbakhsh, M., see Selkirk, M.E., 31

Zarlenga, D.S. and Gamble, H.R.
Molecular cloning and expression of an immunodominant 53-kDa excretory-secretoryantigen from *Trichinella spiralis* muscle larvae, 165

Subject index

Acidification, 257
Alcohol dehydrogenase, 205
Aldehyde reductase, 205
Aldolase, 235
Alpha-amanitin-resistant transcription, 1
Anaerobic-aerobic transition, 63
Antifolate, 21
Antigenic variation, 125
Antigenic variation, 143
Antigenic variation, 285

Brugia malayi, 31

Ca^{2+} mobilization, 101
Ca^{2+} transport, 119
cGMP, 101
Chloroquine accumulation, 257
Chloroquine, 257
Chromosomal alteration, 133
Chromosome, 21
Chromosome size polymorphism, 83
Collagen, 155
Crithidia fasciculata, 119
Crithidia luciliae, 109
Cuticle, 155
Cuticle, 31
Cysteine-rich proteins, 125
Cytoadherence, 189

2-Deoxyglucose, 197
Dihydrofolate reductase-thymidylate synthase point mutation, 83
DNA sequence, 285, 296
Drug resistance, 133
Drug resistance, 21

Eggshell protein, 69
ES antigen, 163
Exflagellation, 101
Evolution, 293

Fermentation product, 63

Gametogenesis, 101
Gene, 241
Gene amplification, 133
Gene cloning, 163
Gene conversion, 293
Gene family, 69
Genetic exchange, 45
Giardia lamblia, 63
Giardia, 125
Glucose metabolism, 109
Glucose transporter, 197
Glutathione transferase, 205
Glycoprotein, 289
Glycosome, 293
Gp29, 31

Haemonchus contortus, 155
Heat shock protein 83, 247
Helminth, 203
Herpetomonas samuelpessoai, 289

Immunity, 241
Ingi, 143
Initiation codon, 55
Intergenic sequence, 247
Intracellular transport, 93

Kinetics, 197
Kinetoplast DNA, 45
Kinetoplastida, 109

D-Lactate, 277
L-Lactate, 277
Leishmania agamae, 119
Leishmania mexicana amazonensis, 247
Leisnmania mexicana, 119
Leishmania mexicana, 269
Leishmania, 225
Lipid peroxidation, 205
Lipophosphoglycan, 225

Malaria, 21, 189, 241, 257
Membrane vesicle, 257
Merozoite antigen, 285
Metacyclogenesis, 225
Methotrexate, 133
Methylglyoxal, 277
Microbody, 293
Minicircle DNA, 175
Mitochondrial DNA, 45
Mitochondrion, 119
Molecular karyotype, 133
Molecular karyotype, 213

Nematode, 31
Nuclear extract, 69

Oligosaccharide, 289

^{31}P magnetic resonance spectroscopy, 109
Parasite diagnosis, 165
Peptidyl aminomethylcoumarin, 269
PfEMP 1, 189
pH, 109
Phosphoglycerate kinase, 293
Plasmodium berghei, 101
Plasmodium falciparum, 55, 83, 93, 101, 189, 235, 257, 277, 285

Plasmodium yoelii, 241
Ploidy, 213
Polymerase chain reaction, 175
Polyphosphate, 109
Procyclic form, 197
Promoter, 1
Proteinase, 269
Proton NMR, 63
Pulsed field gel electrophoresis, 213
Purification, 269
Pyrimethamine resistance, 83

Radioiodination, 155
Recombination, 143
Restriction fragment length polymorphism, 13
Retroposon, 143
Rhoptry, 235
Rhoptry, 241
R_I protein complex, 235
Ring-infected erythrocyte surface antigen, 189

Schistosome, 69
Schizodeme, 175
Secretion, 93
Serine-rich protein, 93
Specificity, 269
Surface antigen, 1, 31, 125
Swine, 165

Taxonomy, 13
Transcription, 69
Translation, 55
Trichinellosis, 165
Trypanosomatid, 289
Trypanosoma brucei, 45, 197, 293
Trypanosoma conorhini rRNA genes, 13
Trypanosoma cruzi rRNA genes, 13
Trypanosoma cruzi, 175, 213
Trypanosome repeated sequence, 143
Trypanosome, 1

UV light transcription mapping, 1

Vaccine, 285